

Referenties

Ontwikkeling van een nieuw meetinstrument voor screening, monitoring en evaluatie in de Nederlandse verslavingszorg

Lisanne Koers, Boukje A.G. Dijkstra, Laura DeFuentes-Merillas, Anneke E. Goudriaan, Vincent M. Hendriks, Jeannette van Manen, Wiebren Markus, Wendy van der Sterren-Kusters, Masha E. Spits, Simon D. Venema, Marleen M. de Waal, Femke Atsma & Arnt F.A. Schellekens

- [1] Degenhardt L, Hall W. Extent of illicit drug use and dependence, and their contribution to the global burden of disease. *Lancet*. 2012;379(9810):55-70.
- [2] Castelpietra G, Knudsen AKS, Agardh EE, Armocida B, Beghi M, Iburg KM, et al. The burden of mental disorders, substance use disorders and self-harm among young people in Europe, 1990-2019: Findings from the Global Burden of Disease Study 2019. *Lancet Reg Health Eur*. 2022;16:100341.
- [3] World Health O. Global status report on alcohol and health and treatment of substance use disorders. World Health Organization; 2024. Report No.: 978-92-4-009674-5.
- [4] Degenhardt L, Charlson F, Ferrari A, Santomauro D, Erskine H, Mantilla-Herrera A, et al. The global burden of disease attributable to alcohol and drug use in 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Psychiatry*. 2018;5(12):987-1012.
- [5] Carroll KM, Onken LS. Behavioral therapies for drug abuse. *Am J Psychiatry*. 2005;162(8):1452-60.
- [6] Volkow ND, Blanco C. Substance use disorders: a comprehensive update of classification, epidemiology, neurobiology, clinical aspects, treatment and prevention. *World Psychiatry*. 2023;22(2):203-29.
- [7] Carroll KM. The profound heterogeneity of substance use disorders: Implications for treatment development. *Curr Dir Psychol Sci*. 2021;30(4):358-64.
- [8] Rutten RJ, Broekman T, van den Brink W, Schippers GM. Differentiating treatment-seeking substance-use disordered patients: support for a staging model. *Sucht*. 2017.
- [9] Myers B, Da Silva N, McLaughlin S, Purnomo J, Shumskaya D, Koume K, et al. The relationship between patient-centred care for substance use disorders and patient outcomes: A scoping review. *Int J Drug Policy*. 2025;139:104770.
- [10] Scott K, Lewis CC. Using measurement-based care to enhance any treatment. *Cognitive and behavioral practice*. 2015;22(1):49-59.
- [11] Hatfield D, McCullough L, Frantz SH, Krieger K. Do we know when our clients get worse? an investigation of therapists' ability to detect negative client change. *Clin Psychol Psychother*. 2010;17(1):25-32.
- [12] Barkham M, De Jong K, Delgado J, Lutz W. Routine Outcome Monitoring (ROM) and Feedback: Research Review and Recommendations. *Psychother Res*. 2023;33(7):841-55.
- [13] Cordony I, Mills L, Mammen K, Lintzeris N. A systematic review on the effect of routine outcome monitoring and feedback on client outcomes in alcohol and other drug treatment. *Drug Alcohol Rev*. 2023;42(7):1701-22.
- [14] de Jong K, Conijn JM, Gallagher RAV, Reshetnikova AS, Heij M, Lutz MC. Using progress feedback to improve outcomes and reduce drop-out, treatment duration, and deterioration: A multilevel meta-analysis. *Clin Psychol Rev*. 2021;85:102002.
- [15] Lappan SN, Brown AW, Hendricks PS. Dropout rates of in-person psychosocial substance use disorder treatments: a systematic review and meta-analysis. *Addiction*. 2020;115(2):201-17.
- [16] Black N, Chung S, Tisdale C, Fialho LS, Aramrattana A, Assanangkornchai S, et al. An International, Multidisciplinary Consensus Set of Patient-Centered Outcome Measures for Substance-Related and Addictive Disorders. *Journal of Clinical Medicine*. 2024;13(7):2154.
- [17] Crunelle CL, van den Brink W, Moggi F, Konstenius M, Franck J, Levin FR, et al. International Consensus Statement on Screening, Diagnosis and Treatment of Substance Use Disorder Patients with Comorbid Attention Deficit/Hyperactivity Disorder. *Eur Addict Res*. 2018;24(1):43-51.
- [18] Patton SC, Watkins LE, Killeen TK, Hien DA. Posttraumatic Stress Disorder and Substance Use Disorder Screening, Assessment, and Treatment. *Curr Psychiatry Rep*. 2024;26(12):843-51.

- [19] Back S, Dansky BS, Coffey SF, Saladin ME, Sonne S, Brady KT. Cocaine dependence with and without post-traumatic stress disorder: a comparison of substance use, trauma history and psychiatric comorbidity. *American Journal on Addictions*. 2000;9(1):51-62.
- [20] Gielen N, Havermans R, Tekelenburg M, Jansen A. Prevalence of post-traumatic stress disorder among patients with substance use disorder: it is higher than clinicians think it is. *European Journal of psychotraumatology*. 2012;3(1):17734.
- [21] Driessen M, Schulte S, Luedecke C, Schaefer I, Sutmann F, Ohlmeier M, et al. Trauma and PTSD in patients with alcohol, drug, or dual dependence: A multi-center study. *Alcoholism: Clinical and Experimental Research*. 2008;32(3):481-8.
- [22] Harrington T, Newman E. The psychometric utility of two self-report measures of PTSD among women substance users. *Addictive behaviors*. 2007;32(12):2788-98.
- [23] Ouimette P, Read J, Brown PJ. Consistency of retrospective reports of DSM-IV Criterion A traumatic stressors among substance use disorder patients. *Journal of Traumatic Stress: Official Publication of The International Society for Traumatic Stress Studies*. 2005;18(1):43-51.
- [24] Reynolds M, Mezey G, Chapman M, Wheeler M, Drummond C, Baldacchino A. Co-morbid post-traumatic stress disorder in a substance misusing clinical population. *Drug and alcohol dependence*. 2005;77(3):251-8.
- [25] Read JP, Brown PJ, Kahler CW. Substance use and posttraumatic stress disorders: Symptom interplay and effects on outcome. *Addictive behaviors*. 2004;29(8):1665-72.
- [26] van Emmerik-van Oortmerssen K, van de Glind G, van den Brink W, Smit F, Crunelle CL, Swets M, et al. Prevalence of attention-deficit hyperactivity disorder in substance use disorder patients: a meta-analysis and meta-regression analysis. *Drug Alcohol Depend*. 2012;122(1-2):11-9.
- [27] Hendriks VM, Kaplan CD, van Limbeek J, Geerlings P. The Addiction Severity Index: reliability and validity in a Dutch addict population. *J Subst Abuse Treat*. 1989;6(2):133-41.
- [28] McLellan AT, Luborsky L, Woody GE, O'Brien CP. An improved diagnostic evaluation instrument for substance abuse patients. The Addiction Severity Index. *J Nerv Ment Dis*. 1980;168(1):26-33.
- [29] Schippers GM, Broekman TG, Buchholz A, Koeter MW, van den Brink W. Measurements in the Addictions for Triage and Evaluation (MATE): an instrument based on the World Health Organization family of international classifications. *Addiction*. 2010;105(5):862-71.
- [30] Verslavingskunde Nederland 2025 [Available from: <https://verslavingskundenederland.nl/>].
- [31] Prinsen CAC, Vohra S, Rose MR, Boers M, Tugwell P, Clarke M, et al. Guideline for selecting outcome measurement instruments for outcomes included in a Core Outcome Set. *BMC/Springer*; 2016.
- [32] Kok G, Gottlieb NH, Peters GJ, Mullen PD, Parcel GS, Ruiter RA, et al. A taxonomy of behaviour change methods: an Intervention Mapping approach. *Health Psychol Rev*. 2016;10(3):297-312.
- [33] Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum*. 2003;49(2):156-63.
- [34] de Beurs E, van Dyck R, Marquenie LA, Lange A, Blonk RWB. De DASS: Een vragenlijst voor het meten van depressie, angst en stress. *Gedragstherapie*. 2001;34(1):35-54.
- [35] Lovibond SH. *Manual for the depression anxiety stress scales*. Sydney psychology foundation. 1995.
- [36] van Dam D, Ehring T, Vedel E, Emmelkamp PM. Validation of the Primary Care Posttraumatic Stress Disorder screening questionnaire (PC-PTSD) in civilian substance use disorder patients. *J Subst Abuse Treat*. 2010;39(2):105-13.
- [37] Kessler RC, Adler L, Ames M, Demler O, Faraone S, Hiripi E, et al. The World Health Organization Adult ADHD Self-Report Scale (ASRS): a short screening scale for use in the general population. *Psychol Med*. 2005;35(2):245-56.
- [38] Neale J, Vitoratou S, Finch E, Lennon P, Mitcheson L, Panebianco D, et al. DEVELOPMENT AND VALIDATION OF 'SURE': A PATIENT REPORTED OUTCOME MEASURE (PROM) FOR RECOVERY FROM DRUG AND ALCOHOL DEPENDENCE. *Drug Alcohol Depend*. 2016;165:159-67.
- [39] Van Nieuwenhuizen C, Janssen-de Ruijter E, Nugter M. Handleiding Manchester short assessment of quality of life (MANSA). 2017.
- [40] van Spijker BA, Batterham PJ, Calear AL, Farrer L, Christensen H, Reynolds J, et al. The suicidal ideation attributes scale (SIDAS): Community-based validation study of a new scale for the measurement of suicidal ideation. *Suicide Life Threat Behav*. 2014;44(4):408-19.
- [41] McLellan AT, Kushner H, Metzger D, Peters R, Smith I, Grissom G, et al. The Fifth Edition of the Addiction Severity Index. *J Subst Abuse Treat*. 1992;9(3):199-213.
- [42] Jellinek. *Utrechtse Somatische Screeningslijst (USS)*. Amsterdam: Jellinek; 2018.

- [43] Bernstein DP, Fink L, Handelsman L, Foote J, Lovejoy M, Wenzel K, et al. Initial reliability and validity of a new retrospective measure of child abuse and neglect. *American Journal of Psychiatry*. 1994;151(8):1132-6.
- [44] Akwa GGZ. Ontwikkelingen COi 2024: Aanvullende thema's en vragen voor ambulante en klinische ggz en verslavingszorg (Versie 6.0). Akwa GGZ; 2024 2024-05.
- [45] Terwee CB, Prinsen C, Chiarotto A, De Vet H, Bouter LM, Alonso J, et al. COSMIN methodology for assessing the content validity of PROMs—user manual. Amsterdam: VU University Medical Center. 2018:1159-70.
- [46] Kirwan JR, Fries JF, Hewlett S, Osborne RH. Patient perspective: choosing or developing instruments. *The Journal of rheumatology*. 2011;38(8):1716-9.
- [47] Meadows KA. Patient-reported outcome measures: an overview. *British journal of community nursing*. 2011;16(3):146-51.
- [48] Wiering B, de Boer D, Delnoij D. Patient involvement in the development of patient-reported outcome measures: a scoping review. *Health Expect*. 2017;20(1):11-23.
- [49] Staniszewska S, Adebajo A, Barber R, Beresford P, Brady LM, Brett J, et al. Developing the evidence base of patient and public involvement in health and social care research: the case for measuring impact. *International Journal of Consumer Studies*. 2011;35(6):628-32.
- [50] Dyer A, Böhnke JR, Curran D, McGrath K, Toner P. A systematic review of quality of life and health-related quality of life as outcomes in substance and behavioural addictions. *Drug Alcohol Rev*. 2023;42(7):1680-700.
- [51] Laudet AB. The case for considering quality of life in addiction research and clinical practice. *Addict Sci Clin Pract*. 2011;6(1):44-55.